## Pre-operative/Pre-procedure Instructions: Adult

Name:	Physician's nan	ne/phone :		
Procedure:		Date:	Time:	
Report to:	Special N	leeds:		
fever), contact you	doctor or surgeon at once	e. Your NIH do	e Clinical Center (such as a control can be contacted by cal sary to postpone your proce	lling the NIH Page
	after midnight the night be very important to assure yo		ery/procedure (no chewing garthe procedure.	um or candy after
disorder, diabetes, currently taking. Toumadin. If you insulin, please take	high blood pressure). Plenis includes aspirin, NSAIE use an inhaler, please bring half of your normal dose. ery/procedure. Unless inst	ease discuss an Os (isuch as Mo ng it with you. If If you use an o	are taking any medicines (for any other drugs/herbal medical otrin, Aleve) and blood thinned by you use have diabetes melloral pill/hypoglycemic agent, e, take all your medicines as	ations that you are ers (isuch as litus and use please hold it on
A responsible adult	(18 years or older) must a	ccompany you	when you are discharged fro	om the hospital.
Do not drink alcoho	ic beverages for 24 hours	prior to surgery	//procedure.	
Please remove all j valuables at home.		/ piercing jewel	ry) before coming to the hos	pital. Leave all
Please remove you	r make-up before coming	to the Clinical C	Center.	
	prosthetic devices. Thes		if you wear contact lenses, he removed before surgery.	
Expect to remain 1	to 3 hours at the Clinical	Center after the	surgery/procedure.	
		•	Clinical Center. This is unlike ain the reasons for a hospita	•



This information is prepared specifically for patients participating in clinical research at the Warren Grant Magnuson Clinical Center at the National Institutes of Health and is not necessarily applicable to individuals who are patients elsewhere. If you have questions about the information presented here, talk to a member of your healthcare team.

Questions about the Clinical Center? OCCC@cc.nih.gov

Where applicable, brand names of commercial products are provided only as illustrative examples of acceptable products, and do not imply endorsement by NIH; nor does the fact that a particular brand name product is not identified imply that such product is unsatisfactory.

National Institutes of Health Warren Grant Magnuson Clinical Center Bethesda, Maryland 20892